

HIV, HCV and STD Screening Resource Guide

	HIV	HCV	STD
SEX:			
• Sexual partners of HIV infected persons	Annually /3-6 months	At Least Once in Lifetime	Annually/ 3-6months
• Multiple sex partners	Annually		Annually
• Men who have sex with men (MSM)	Annually		Annually /3-6 months
• Having unprotected sex	Annually		Annually
• Unprotected anal sex	Annually		Rectal / Pharyngeal testing
• Persons who exchange sex for money or drugs	Annually /3-6 months	Annually	Annually
• All pregnant women	Test With Each Pregnancy		Test With Each Pregnancy
Drugs:			
• Used drugs not prescribed to them (non-injection)		Test	
• Current or past Injection drug users	Annually /3-6 months	Annually	
• Share or have shared needles or have had needle exposure	Annually	Annually	
Tattoos:			
• Tattoos or body piercings in an unsterile environment	Test At Every New Exposure & At 6 Months Post-Exposure	Test At Every New Exposure & At 6 Months Post-Exposure	
Diagnosed with Disease:			
• Have HIV or AIDS		At Diagnosis/As Needed	At Diagnosis/Annually
• Persons diagnosed with Tuberculosis	At Diagnosis		
• Persons diagnosed with HCV	At Diagnosis		
• Persons diagnosed with a STD	At Diagnosis		
Received Treatment:			
• Received donated blood or organs before 1992		At Least Once	
• Received blood clotting factors before 1987		At Least Once	
• Are getting long term hemodialysis		At Least Once	
• Undiagnosed liver problems		At Least Once	